Amendments to the Claims

This listing of claims will replace all prior versions, and listings, of claims in the application:

Claim 1 (Currently amended): A compound of formula III:

or a pharmaceutically acceptable derivative or prodrug thereof, wherein:

 Z^1 is nitrogen or CR^8 , Z^2 is nitrogen or CH, and Z^3 is nitrogen or CR^x , provided that when one of Z^1 and or Z^3 is nitrogen, the other of Z^1 or Z^3 is CR^8 or CR^x , respectively;

 R^x is T- R^3 or L-Z- R^3 ;

Q is selected from $-N(R^4)$ -, -O-, -S-, or $-CH(R^6)$ -;

R1 is T-(Ring D);

Ring D is a 5-7 membered monocyclic ring or 8-10 membered bicyclic ring selected from aryl, heteroaryl, heterocyclyl or carbocyclyl, said heteroaryl or heterocyclyl ring having 1-4 ring heteroatoms selected from nitrogen, oxygen or sulfur, wherein each substitutable ring carbon of Ring D is independently substituted by oxo, T-R⁵, or V-Z-R⁵, and each substitutable ring nitrogen of Ring D is independently substituted by -R⁴;

T is a valence bond or a C₁₋₄ alkylidene chain, wherein when Q is -CH(R⁶)-, a methylene unit of said C₁₋₄ alkylidene chain is optionally replaced by -O-, -S-, -N(R⁴)-, -CO-, -OC(O)NH-, or -NHCO₂-;

Z is a C₁₋₄ alkylidene chain;

L is -O-, -S-, -SO-, -SO₂-, -N(R⁶)SO₂-, -SO₂N(R⁶)-, -N(R⁶)-, -CO-, -CO₂-, -N(R⁶)CO-, -N(R⁶)C(O)O-, -N(R⁶)CON(R⁶)-, -N(R⁶)SO₂N(R⁶)-, -N(R⁶)N(R⁶)-, -C(O)N(R⁶)-, -C(R⁶)₂SO-, -C(R⁶)₂SO-, -C(R⁶)₂SO₂-, -C(R⁶)₂SO₂N(R⁶)-,

- $-C(R^6)_2N(R^6)-, -C(R^6)_2N(R^6)C(O)-, -C(R^6)_2N(R^6)C(O)O-, -C(R^6)=NN(R^6)-, -C(R^6)=N-O-, -C(R^6)_2N(R^6)N(R^6)-, -C(R^6)_2N(R^6)SO_2N(R^6)-, or -C(R^6)_2N(R^6)CON(R^6)-;$
- R² and R² are independently selected from -R, -T-W-R⁶, or R² and R² are taken together with their intervening atoms to form a fused, 5-8 membered, unsaturated or partially unsaturated, ring having 0-3 ring heteroatoms selected from nitrogen, oxygen, or sulfur, wherein each substitutable ring carbon of said fused ring formed by R² and R² is independently substituted by halo, oxo, -CN, -NO₂, -R⁷, or -V-R⁶, and each substitutable ring nitrogen of said ring formed by R² and R² is independently substituted by R⁴;
- R³ is selected from -R, -halo, -OR, -C(=O)R, -CO₂R, -COCOR, -COCH₂COR, -NO₂, -CN, -S(O)R, -S(O)₂R, -SR, -N(R⁴)₂, -CON(R⁷)₂, -SO₂N(R⁷)₂, -OC(=O)R, -N(R⁷)COR, -N(R⁷)CO₂(C₁₋₆ aliphatic), -N(R⁴)N(R⁴)₂, -C=NN(R⁴)₂, -C=N-OR, -N(R⁷)CON(R⁷)₂, -N(R⁷)SO₂N(R⁷)₂, -N(R⁴)SO₂R, or -OC(=O)N(R⁷)₂;
- each R is independently selected from hydrogen or an optionally substituted group selected from C_{1-6} aliphatic, C_{6-10} aryl, a heteroaryl ring having 5-10 ring atoms, or a heterocyclyl ring having 5-10 ring atoms;
- each R^4 is independently selected from $-R^7$, $-CO_2$ (optionally substituted C_{1-6} aliphatic), $-CON(R^7)_2$, or $-SO_2R^7$;
- each R^5 is independently selected from -R, halo, -OR, -C(=O)R, -CO₂R, -COCOR, -NO₂, -CN, -S(O)R, -SO₂R, -SR, -N(R^4)₂, -CON(R^4)₂, -SO₂N(R^4)₂, -OC(=O)R, -N(R^4)COR, -N(R^4)CO₂(optionally substituted C₁₋₆ aliphatic), -N(R^4)N(R^4)₂, -C=NN(R^4)₂, -C=N-OR, -N(R^4)CON(R^4)₂, -N(R^4)SO₂N(R^4)₂, -N(R^4)SO₂R, or -OC(=O)N(R^4)₂;
- V is -O-, -S-, -SO-, -SO₂-, -N(R⁶)SO₂-, -SO₂N(R⁶)-, -N(R⁶)-, -CO-, -CO₂-, -N(R⁶)CO-,
 -N(R⁶)C(O)O-, -N(R⁶)CON(R⁶)-, -N(R⁶)SO₂N(R⁶)-, -N(R⁶)N(R⁶)-, -C(O)N(R⁶)-,
 -OC(O)N(R⁶)-, -C(R⁶)₂O-, -C(R⁶)₂S-, -C(R⁶)₂SO-, -C(R⁶)₂SO₂-, -C(R⁶)₂SO₂N(R⁶)-,
 -C(R⁶)₂N(R⁶)-, -C(R⁶)₂N(R⁶)C(O)-, -C(R⁶)₂N(R⁶)C(O)O-, -C(R⁶)=NN(R⁶)-,
 -C(R⁶)₂N(R⁶)N(R⁶)-, -C(R⁶)₂N(R⁶)SO₂N(R⁶)-, or -C(R⁶)₂N(R⁶)CON(R⁶)-;
- W is $-C(R^6)_2O_-$, $-C(R^6)_2S_-$, $-C(R^6)_2SO_-$, $-C(R^6)_2SO_2$, $-C(R^6)_2SO_2N(R^6)_-$, $-C(R^6)_2N(R^6)_-$, or $-C(R^6)_-$;

01/18/2005 10:53 FAX @005/010

each R⁶ is independently selected from hydrogen or an optionally substituted C₁₋₄ aliphatic group, or two R⁶ groups on the same nitrogen atom are taken together with the nitrogen atom to form a 5-6 membered heterocyclyl or heteroaryl ring;

- each R⁷ is independently selected from hydrogen or an optionally substituted C₁₋₆ aliphatic group, or two R⁷ on the same nitrogen are taken together with the nitrogen to form a 5-8 membered heterocyclyl or heteroaryl ring; and
- R^8 is selected from -R, halo, -OR, -C(=O)R, -CO₂R, -COCOR, -NO₂, -CN, -S(O)R, -SO₂R, -SR, -N(R⁴)₂, -CON(R⁴)₂, -SO₂N(R⁴)₂, -OC(=O)R, -N(R⁴)COR, -N(R⁴)CO₂(optionally substituted C₁₋₆ aliphatic), -N(R⁴)N(R⁴)₂, -C=NN(R⁴)₂, -C=N-OR, -N(R⁴)CON(R⁴)₂, -N(R⁴)SO₂R, or -OC(=O)N(R⁴)₂.

Claim 2 (Currently amended): The compound according to claim 1, wherein Q is $-N(R^4)$ -, -S-, or $-CH(R^6)$ -, and said compound is of formula ΠIa , or ΠIb , HIe, or IIId:

or a pharmaceutically acceptable derivative or prodrug thereof.

Claim 3 (Original): The compound according to claim 2, wherein said compound has one or more features selected from the group consisting of:

- (a) Rx is hydrogen, alkyl- or dialkylamino, acetamido, or a C1-4 aliphatic group;
- (b) R¹ is T-(Ring D), wherein T is a valence bond or a methylene unit;
- (c) Ring D is a 5-7 membered monocyclic or an 8-10 membered bicyclic aryl or heteroaryl ring; and
- (d) R² is -R or -T-W-R⁶ and R² is hydrogen, or R² and R² are taken together to form an optionally substituted benzo ring.

Claim 4 (Original): The compound according to claim 3, wherein:

- (a) R^x is hydrogen, alkyl- or dialkylamino, acetamido, or a C₁₋₄ aliphatic group;
- (b) R¹ is T-(Ring D), wherein T is a valence bond or a methylene unit;
- (c) Ring D is a 5-7 membered monocyclic or an 8-10 membered bicyclic aryl or heteroaryl ring; and
- (d) R² is -R or -T-W-R⁶ and R² is hydrogen, or R² and R² are taken together to form an optionally substituted benzo ring.

Claim 5 (Original): The compound according to claim 3, wherein said compound has one or more features selected from the group consisting of:

- (a) R¹ is T-(Ring D), wherein T is a valence bond, and Q is -S- or -NH-;
- (b) Ring D is a 5-6 membered monocyclic or an 8-10 membered bicyclic aryl or heteroaryl ring; and
- (c) R² is -R and R² is hydrogen, wherein R is selected from hydrogen, C₁₋₆ aliphatic, phenyl, a 5-6 membered heteroaryl ring, or a 5-6 membered heterocyclic ring.

Claim 6 (Original): The compound according to claim 5, wherein:

- (a) R¹ is T-(Ring D), wherein T is a valence bond, and Q is -S- or -NH-;
- (b) Ring D is a 5-6 membered monocyclic or an 8-10 membered bicyclic aryl or heteroaryl ring; and
- (c) R² is -R and R² is hydrogen, wherein R is selected from hydrogen, C_{1.6} aliphatic, phenyl, a 5-6 membered heteroaryl ring, or a 5-6 membered heterocyclic ring.

Claim 7 (Original): The compound according to claim 5, wherein said compound has one or more features selected from the group consisting of:

- (a) R^x is hydrogen methyl, ethyl, propyl, cyclopropyl, isopropyl, methylamino or acetamido;
- (b) R¹ is T-(Ring D), wherein T is a valence bond and Ring D is a 5-6 membered aryl or heteroaryl ring, wherein Ring D is optionally substituted with one to two groups selected from -halo, -CN, -NO₂, -N(R⁴)₂, optionally substituted C₁₋₆ aliphatic group,

-OR, -CO₂R, -CON(R^4)₂, -OCO(R^4)₂, -N(R^4)COR, -N(R^4)SO₂R, -N(R^6)COCH₂CH₂N(R^4)₂, or -N(R^6)COCH₂CH₂CH₂N(R^4)₂; and

(c) R² is hydrogen or a substituted or unsubstituted C₁₋₆ aliphatic.

Claim 8 (Original): The compound according to claim 7, wherein:

- (a) R^x is hydrogen methyl, ethyl, propyl, cyclopropyl, isopropyl, methylamino or acetamido;
- (b) R¹ is T-(Ring D), wherein T is a valence bond and Ring D is a 5-6 membered aryl or heteroaryl ring, wherein Ring D is optionally substituted with one to two groups selected from -halo, -CN, -NO₂, -N(R⁴)₂, optionally substituted C₁₋₆ aliphatic group, -OR, -CO₂R, -CON(R⁴)₂, -OCO(R⁴)₂, -N(R⁴)COR, -N(R⁴)SO₂R, -N(R⁶)COCH₂CH₂N(R⁴)₂, or -N(R⁶)COCH₂CH₂N(R⁴)₂; and
- (c) R² is hydrogen or a substituted or unsubstituted C₁₋₆ aliphatic.

Claim 9 (Currently amended): A compound selected from the group consisting of:

N⁵-(1H-Indazol 6 yl) N³ (5 methyl 1H pyrazol 3 yl) [1,2,4]triazine 3,5 diamine;

N-[4-[3-(5 Methyl 1H pyrazol 3 ylamino) [1,2,4]triazin 5 ylsulfanyl] phenyl) acetamide;

[5-(3 Methoxy benzyl) [1,2,4]triazin 3 yl] (5 methyl 1H pyrazol 3 yl) amine;

N³-(5-Cyclopropyl 1H pyrazol 3 yl) N⁵-pyridin 3 ylmethyl [1,2,4]triazine 3,5 diamine;

[5-(Benzothiazol 6 ylsulfanyl) [1,2,4]triazin 3 yl] (5 cyclopropyl 1H pyrazol 3 yl) amine;

[4-[3-(5 Cyclopropyl 1H pyrazol 3 ylamino) [1,2,4]triazin 5 yloxy] phenyl) acetonitrile;

N-[4-[3-(1H Indazol 3 ylamino) [1,2,4]triazin 5 ylamino] phenyl) methanesulfonamide;

(1H Indazol 3 yl) [5-(thiophen 2 ylmethylsulfanyl) [1,2,4]triazin 3 yl] amine;

N⁵-(5 Methyl 1H pyrazol 3 yl) N³ pyridin 3 ylmethyl [1,2,4]triazine 3,5 diamine;

[3-(Benzothiazol 6 ylsulfanyl) [1,2,4]triazin 5 yl] (5 methyl 1H pyrazol 3 yl) amine;

[4-[5-(5 Methyl 1H pyrazol 3 ylamino) [1,2,4]triazin 3 yloxy] phenyl) acetonitrile;

N⁵-(5 Cyclopropyl 1H pyrazol 3 ylamino) [1,2,4]triazin 3 ylsulfanyl] phenyl) acetomide:

N⁵-(1*H*-Indazol 3-yl) N³-(1*H* indazol 6-yl) [1,2,4]triazine-3,5-diamine; (1*H* Indazol 3-yl) [3 (3-methoxy-phenylsulfanyl) [1,2,4]triazin-5-yl] amine; N⁵-(1*H*-Indazol-6-yl)-N³-(5-methyl-1*H*-pyrazol-3-yl)-pyridazine-3,5-diamine; N-{4-[6-(5-Methyl-1*H*-pyrazol-3-ylamino)-pyridazin-4-ylsulfanyl]-phenyl}-acetamide; [5-(3-Methoxy-benzyl)-pyridazin-3-yl]-(5-methyl-1*H*-pyrazol-3-yl)-amine; N³-(5-Cyclopropyl-1*H*-pyrazol-3-yl)-N⁵-pyridin-3-ylmethyl-pyridazine-3,5-diamine; [5-(Benzothiazol-6-ylsulfanyl)-pyridazin-3-yl]-(5-cyclopropyl-1*H*-pyrazol-3-yl)-amine; {4-[6-(5-Cyclopropyl-1*H*-pyrazol-3-ylamino)-pyridazin-4-yloxy]-phenyl}-acetonitrile; N-{4-[6-(1*H*-Indazol-3-ylamino)-pyridazin-4-ylamino]-phenyl}-methanesulfonamide; (1*H*-Indazol-3-yl)-[5-(thiophen-2-ylmethylsulfanyl)-pyridazin-3-yl]-amine; N⁵-(5-Methyl-1*H*-pyrazol-3-yl)-N³-pyridin-3-ylmethyl-pyridazine-3,5-diamine; [6-(Benzothiazol-6-ylsulfanyl)-pyridazin-4-yl]-(5-methyl-1*H*-pyrazol-3-yl)-amine; {4-[5-(5-Methyl-1*H*-pyrazol-3-ylamino)-pyridazin-3-yloxy]-phenyl}-acetonitrile; N⁵-(5-Cyclopropyl-1*H*-pyrazol-3-yl)-N³-(1*H*-indazol-6-yl)-pyridazine-3,5-diamine; N-{4-[5-(5-Cyclopropyl-1*H*-pyrazol-3-ylamino)-pyridazin-3-ylsulfanyl]-phenyl}-acetamide; N⁵-(1*H*-Indazol-3-yl)-N³-(1*H*-indazol-6-yl)-pyridazine-3,5-diamine; and (1*H*-Indazol-3-yl)-[6-(3-methoxy-phenylsulfanyl)-pyridazin-4-yl]-amine.

Claim 10 (Original): A composition comprising a compound according to any of claims 1-9, and a pharmaceutically acceptable carrier.

Claim 11 (Original): The composition according to claim 10, further comprising an additional therapeutic agent.

Claim 12 (Original): A method of inhibiting Aurora-2 or GSK-3 activity in a biological sample comprising the step of contacting said biological sample with a compound according to any one of claims 1-9.

Claim 13 (Original): A method of inhibiting Aurora-2 activity in a patient comprising the step of administering to said patient a composition according to claim 10.

Claim 14 (Original): A method of inhibiting Aurora-2 activity in a patient comprising the step of administering to said patient a composition according to claim 11.

Claim 15 (Original): A method of treating an Aurora-2-mediated disease, which method

comprises administering to a patient in need of such a treatment a therapeutically effective amount of a composition according to claim 10.

Claim 16 (Original): The method according to claim 15, wherein said disease is selected from colon, breast, stomach, or ovarian cancer.

Claim 17 (Original): The method according to claim 16, wherein said method further comprises administering an additional therapeutic agent.

Claim 18 (Original): The method according to claim 17, wherein said additional therapeutic agent is a chemotherapeutic agent.

Claim 19 (Original): A method of inhibiting GSK-3 activity in a patient comprising the step of administering to said patient a composition according to claim 10.

Claim 20 (Original): A method of inhibiting GSK-3 activity in a patient comprising the step of administering to said patient a composition according to claim 11.

Claim 21 (Original): A method of method of treating a GSK-3-mediated disease, which method comprises administering to a patient in need of such a treatment a therapeutically effective amount of a composition according to claim 10.

Claim 22 (Original): The method according to claim 21, wherein said GSK-3-mediated disease is selected from diabetes, Alzheimer's disease, Huntington's Disease, Parkinson's Disease, AIDS-associated dementia, amyotrophic lateral sclerosis (AML), multiple sclerosis (MS), schizophrenia, cardiomycete hypertrophy, reperfusion/ischemia, or baldness.

Claim 23 (Original): The method according to claim 22, wherein said GSK-3-mediated disease is diabetes.

01/18/2005 10:55 FAX **2010/010**

Claim 24 (Original): A method of enhancing glycogen synthesis or lowering blood levels of

glucose in a patient in need thereof, which method comprises administering to said patient a

therapeutically effective amount of a composition according to claim 10.

Claim 25 (Original): A method of inhibiting the production of hyperphosphorylated Tau protein

in a patient, which method comprises administering to a patient in need thereof a therapeutically

effective amount of a composition according to claim 10.

Claim 26 (Original): A method of inhibiting the phosphorylation of \(\beta \)-catenin, which method

comprises administering to a patient in need thereof a therapeutically effective amount of a

composition according to claim 10.

Applicants request entry of the above amendments, favorable consideration of the

application, and early allowance of the pending claims.

Respectfully submitted,

Lisa Dixon Reg. No. 40,995

Attorney for Applicants

VERTEX PHARMACEUTICALS INCORPORATED

130 Waverly Street

Cambridge, Massachusetts 02139

Tel.: (617) 444-6396

Fax.: (617) 444-6438

9